

Landowners Guide for Controlling Phragmites

An aggressive, nonnative variety of phragmites (*Phragmites australis*), also known as common reed, is threatening the ecological health of Lancaster County wetlands and riparian areas.



Phragmites with mature seed heads

This invasive variety of phragmites is becoming widespread throughout Lancaster County. Even though the infested acres are quite small, infestations are scattered throughout the county and have enough seed production to potentially infest any areas having a saturated soil condition sometime during the year. The infestations have been found along streams, wetlands, ponds, lagoons, road ditches and railroads. Once started in these areas, it quickly forms a colony that completely crowds out all other plants including cattails. If these scattered infestations are not controlled, solid stands of phragmites will completely cover wetlands and the riparian areas along upland streams and Salt Creek. The rapid expansion of this variety of phragmites will result in adverse ecological, economic and social impacts on the natural resources, the people of Lancaster County and the entire Lower Platte River Basin.

The easiest way to control phragmites is to begin a control program as soon as it is observed on your property, before the plants become well established. With the cooperation of landowners, complete eradication of this early invasion of phragmites may be achievable. An aggressive approach is needed now to prevent the rapid expansion to all Lancaster County riparian areas including Salt Creek.

Contact our office to report an infestation and / or for assistance in developing a control plan.

Lancaster County Weed Control Authority
444 Cherry Creek Rd, Building B, Lincoln, Nebraska 68528
402-441-7817

Brent Meyer, Superintendent
bmeyer@lancaster.ne.gov
www.lancaster.ne.gov/weed

Understanding Phragmites

To better control and manage phragmites it is helpful to understand the physical characteristics of the plant, as well as how and when it reproduces and spreads. In Nebraska, phragmites is growing along rivers, streams, flood plains and lakeshores with heaviest concentrations in the central Platte area but is being found throughout the Platte River and other rivers. It now is also beginning to be found in ditches and other low wet areas. Phragmites continues to expand within Nebraska, in part because it reproduces through wind and water dispersal of seeds and vigorous vegetative reproduction through rhizomes which can grow 30 feet or more in one year. Rhizomes



Phragmites stolons

Phragmites turns a tan color in the fall and most leaves drop off, leaving only the stalk and plume-topped shoot commonly seen throughout winter. Each mature plant can produce as many as 2,000 seeds annually. New stands of phragmites may develop from seed which is spread by water and the wind, although this is a slower process than spreading by rhizome fragments.

Currently, native phragmites has not been identified in Lancaster County. However, it may be present, so it is important to identify the native phragmites versus the non-native invasive variety before attempting control.

Additional information on how to identify native versus non-native phragmites can be found at <http://www.invasiveplants.net/phragmites/phrag/morph.htm>

Contact our office and we will make the identification and provide assistance in developing a control plan.

Recommended Management Strategies

Controlling the spread of phragmites is crucial to the restoration of native wetland plant communities and protection of vital fish and wildlife habitat. Phragmites can easily spread if improper control methods are used. The following guidelines will help ensure that phragmites control efforts are effective.

Phragmites can be controlled using an initial herbicide treatment followed by mechanical removal (e.g., cutting, mowing) and annual maintenance. *(It is recommended to wait for at least 12 months after herbicide application to mow or burn)* For large areas with dense stands of phragmites, prescribed burning used after herbicide treatment can provide additional control and ecological benefits over mechanical removal. However, phragmites burns very hot and fast, and prescribed burns should be performed only by trained personnel. A burn permit and approval is required from local fire departments. No biological control methods for phragmites are currently available. However, researchers at Cornell University are studying several insects native to Europe that are known to attack phragmites as possible biological controls. To date, field experience and research have shown that using herbicides is the most effective method and is recommended as the first step toward effective control of phragmites.

Glyphosate (Rodeo) and **imazapyr** (Arsenal or Habitat) are two herbicides known to be effective in controlling phragmites. These herbicides are non-selective and will affect any plant species through contact with the leaves and stems. However, when applied using the correct method and used according to chemical manufacturer's instructions, impacts to native plants, as well as mammals, birds, and fish can be minimized. The aquatic formulations of these herbicides are required for use in wetlands. A surfactant should be added to these aquatic formulations to improve the effectiveness of the treatment.

Herbicide Selection

While the cost per gallon of Arsenal or Habitat (imazapyr) can be higher than Rodeo or other glyphosate products, results from recent studies suggest that imazapyr used alone or in combination with glyphosate can control phragmites for a longer period of time. When using herbicides, phragmites should be treated in early to late summer (June – September) using imazapyr, or late summer (August – September) using either glyphosate or a glyphosate / imazapyr mixture, to achieve effective control. Indications are that earlier applications are the most effective.

Methods of Application

Numerous methods may be used to apply these herbicides, depending on the size of the phragmites stand and existing site conditions.

Herbicide application methods for scattered plants or isolated plant stands include: injecting stems, hand swiping or selective hand spraying.



Seed head before maturity

Spot treating areas with scattered plants or isolated stands can prevent the establishment of large dense stands and is more cost effective.

Large dense stands may require use of commercial equipment. The use of a licensed or certified applicator is recommended to minimize damage to native plants and to ensure that safety requirements are met. The use of a licensed applicator certified in aquatic pest management is recommended for herbicide application in wetlands. Commercial pesticide use certification is required prior to using Habitat according to the manufacturer's label and is recommended prior to using Arsenal or glyphosate.

As with most invasive plants and animals, complete eradication of phragmites is unlikely.

Phragmites control requires a commitment to an integrated and long-term management approach. To achieve desired results, herbicides must be used in conjunction with mechanical methods or burning, and re-applied in subsequent years to spot-treat individual plants or patches of plants that were not completely eliminated in the first application. Large, dense phragmites stands will likely require follow-up spot treatments, and phragmites will continue to re-establish from remnant and neighboring populations, as well as the existing seed bank.

Phragmites typically begins to recover two to three years after treatment and will re-establish unless follow-up annual maintenance occurs, including spot treatment with herbicides.

Application Rates

Wet Sites - Sites with standing water at the time of the application must use an aquatic labeled herbicide. Recommendations are based on information from the UNL Extension EC 130 Guide for Weed Management in Nebraska. Always read and follow labels directions.

ARSENAL / HABITAT (Imazapyr) Apply to actively growing green foliage after full leaf elongation and up to first killing frost, ensure 100% coverage. (i.e., July up to first killing frost) Earlier applications have shown to be most effective. If the stand has a substantial amount of old stem tissue you may need mow or burn and allow to regrow to approximately 5' tall before treatment.

Application rates for Imazapyr – 2.0 – 4.0 pints per acre. Always use a label recommended surfactant.

AQUATIC GLYPHOSATE - Apply after plants are in full bloom and up to first killing frost, ensure 100% coverage. (i.e., Late August up to first killing frost). Earlier applications have shown to be most effective. If the stand has a substantial amount of old stem tissue you may need to mow or burn and allow to regrow to approximately 5' tall before treatment.

Application rates for glyphosate – 3.0 – 4.0 quarts per acre. Always use a label recommended surfactant.


Commercial Applicators

We have a list of commercial applicators including aerial applicators that would apply herbicides for the control of phragmites. Contact our office for a list or other assistance in developing a control plan.

Lancaster County Weed Control Authority
444 Cherry Creek Rd, Building B, Lincoln, Nebraska 68528
402-441-7817

Brent Meyer, Superintendent
bmeyer@lancaster.ne.gov
www.lancaster.ne.gov/weeds

2011 Phargmites



map: Lancaster County Weed Control
November 28, 2011